

Rob Hannon - Direct Access Endoscopy - Referral Form

Fax to:

email to:

PATIENT DETAILS

Name:

Date of Birth:

Address:

Tel:

Mob:

GENERAL PRACTITIONER DETAILS

Referring GP Name:

Address:

Tel:

Mob:

INVESTIGATION REQUIRED & PRESENTING PROBLEM

Colonoscopy

Rectal Bleeding

Altered bowel habit

Lower Abdominal pain

Anaemia

Family hx

Weight Loss

Gastroscopy

Dysphagia

Dyspepsia (despite PPI)

Upper Abdominal pain

Anaemia

Gastroscopy & Colonoscopy

PAST MEDICAL HISTORY

A.Fib.

CVA

IHD

IDDM/NIDDM

MEDICATION

Is the patient taking:

Aspirin

Plavix

Warfarin

Dabigatran/ Rivaroxaban (Pradaxa/Xarelto)

Insulin

Oral Hypoglycaemics

Notes:

1. Patients on Aspirin alone do not require to stop taking Aspirin before test.
2. Patients Plavix should discontinue medication for 7 days before the test (after discussion with Cardiologist).

3. Patients on Warfarin should discontinue this Medication 5 days before test & require INR the day before test. They may require bridging LMW Heparin (CVA/ cardiac valve replacement).
4. Patients on novel anticoagulants (Pradaxa/ Xarelto) must stop these 72 hours before test.
5. Diabetic patients are scheduled 1st on the list.
6. Patients will be sent a prescription for **Moviprep (with instructions for use)** prior to the test.